


Authored by
Professor Mark O'Donnell,
Consultant Vascular and
Endovascular Surgeon



Vascular

Foam Sclerotherapy for Varicose Veins

Post-Operative Care Information



KINGSBRIDGE

Private Hospital Group
★★★★★



Belfast | Ballykelly | Newry

kingsbridgeprivatehospital.com



Professor Mark E. O'Donnell
Consultant Vascular and Endovascular Surgeon

Many thanks for attending for your Foam Sclerotherapy Procedure. It is important to take the necessary steps to ensure a smooth and successful recovery. To help guide you through this process, Professor O'Donnell and his varicose vein team have created a post-procedure care leaflet to provide you with important information and instructions to optimise your outcome.

What is Sclerotherapy?

Sclerotherapy is a minimally invasive treatment for varicose veins that involves injecting a foam-based solution directly into the affected vein using a very small needle which causes irritation and inflammation in the targeted blood vessels, leading to their closure and eventual absorption by the body.

Whilst sclerotherapy is effective in improving the appearance of varicose and spider veins, it does not prevent the formation of new ones and often more than one treatment is required as sclerotherapy may not eliminate all veins during a single treatment session, particularly if you have an extensive or complex venous presentation. Multiple treatment sessions may be necessary to achieve the desired results.

After the procedure it is important to take good care of yourself to promote healing, reduce the risk of complications and get a great result.

Post-Operative Care

Here are some tips for sclerotherapy aftercare. Please take time to read through the instructions carefully and if you have any post-operative questions, please contact your clinic; details are provided on pages 8-9.

Prescribed Medications & Pain Medications

Please continue to take all prescribed medications including blood thinners. Professor O'Donnell is aware that certain cardiovascular medications can lead to increased bruising effects. However, it is more important to ensure that your heart health is protected during your procedure. You do not need to fast prior to your procedure and a well-deserved refreshment awaits after your procedure.

Sclerotherapy may cause some discomfort, stinging or itching during the procedure, which normally settles after a few minutes – usually by the time the compression stockings have been applied. It is ok to take some paracetamol to help manage your discomfort if required for the following few days after your procedure. Be sure to follow the instructions and only take medication as directed.

Physical Activity & Walking

You should avoid heavy lifting, strenuous exercise and other activities that may put pressure on your legs for two weeks after the procedure. No gym, not even an upper body work-out, cycling, swimming, golf, yoga, pilates etc. for 2 weeks until the stockings are no longer required.

Walking is a low-impact exercise that can help improve circulation and promote healing. You should try to walk regularly after the procedure, but avoid standing or sitting for long periods of time. Gradually build up your walking over the first few days after your procedure.

Compression Stockings & Leg Dressings

Occasionally small adherent dressings are applied to your lower extremities after foam sclerotherapy. These often fall off in the shower or peel off thereafter when wet. If still adherent to the skin after 7 days, you can carefully remove these dressings yourself and bin them.

Professor O'Donnell recommends that you wear the compression stockings provided for 2 weeks after the procedure. These stockings need to remain in place for the first 48 hours - do not remove them within this period. After 48 hours you may remove the stockings for a shower (short duration and not too warm). You can then reapply the stockings afterwards and shower as often as you like from then on. Avoid hot baths, hot-tubs, saunas or any excessive heat sources to the treated lower extremities for 2 weeks post-procedure until stocking use is no longer required. The stockings help to promote blood flow and prevent blood clots whilst minimising bruising to optimise your procedural result.

Sometimes an additional layer of compression bandaging called "Peha-Haft" is applied on top of your compression stockings. This bandage is self-adherent and can be easily peeled off by you after 48-hours by gradually unwinding the different layers. The "Peha-Haft" bandage can then be binned. The compression stockings can then be easily removed by yourself and kept to the side to be re-applied after you have finished showering.

The stockings are to be worn Day and Night for the first week and only during the daytime for the second week (can be removed at night when sleeping). Please note it is common for stockings to occasionally slip down as they loosen with continued wear. You may need to pull them up again occasionally but if this is persistent you may need a new pair or a smaller size. Please contact the clinic if required.



Patient Expectations & Post-Procedure Healing Process

All patients are different with variations in size and severity of surface area affected by their lower extremity varicose veins. Professor O'Donnell will provide you with an honest assessment of treatment coverage after your procedure where additional sclerotherapy procedures are sometimes/often required to optimise outcomes and treatment completion.

Most patients are nervous prior to and during their procedure. Professor O'Donnell will always try to provide high-quality care in a very informal atmosphere. Most people leave after their sclerotherapy without any discomfort as the foam sclerotherapy usually only causes some stinging and itch for the first few minutes. After two weeks when the compression stockings are removed, it is normal to see a lot of bruising and larger veins may be lumpy. Please do not be alarmed as this is what the foam sclerotherapy procedure is designed to do.

At this stage, if you do have a lot of bruising, it is recommended to massage Arnica cream directly onto the bruises which acts to help reduce bruising (Arnica cream can be purchased in your local pharmacy or via online sources). General moisturiser such as E45 cream can be massaged vigorously (as if "kneading bread") on to the affected sites that are lumpy. You should aim to massage affected areas for 10-15 minutes every day.

It is not uncommon for the treated veins to throb after a few weeks as the sclerotherapy treatment effect evolves. Please do not worry about this which can sometimes reduce your ability to vigorously massage the treated areas. Reduce pressure for a period of time and gradually increase pressure applied as discomfort settles. If this inflammation effect is causing continued discomfort and if it is safe for you to take oral anti-inflammatory medication, Professor O'Donnell recommends that you take oral anti-inflammatory medication such as Ibuprofen 200mg 3 times a day (with meals to avoid irritation to your stomach) for 1-2 weeks. Alternatively you could use Ibuprofen gel on the affected areas (it is important you do not take oral Ibuprofen and use Ibuprofen gel together, select one method only). Both of these medications can be obtained over the counter from your pharmacist.

Bruises and lumps can persist for many months after your initial procedure. Please try not to worry about this. Professor O'Donnell recommends continued massage with moisturiser and Arnica cream as you will continue to heal and results will continue to improve in the 12 months after your procedure. However, most patients see significant results and satisfactory healing progress by 3 months where Professor O'Donnell will usually have you appointed to a scheduled review consultation.

Complications

Although complications are rare, Professor O'Donnell has documented a number of potential concerns below. It's important to watch for signs of infection, bleeding or other problems. If you notice any unusual symptoms, please contact your clinic; details are provided on pages 8-9.

Altered Sensation & Numbness

Certain treatment areas in the lower extremity can develop altered sensations after sclerotherapy in particular numbness. This is more common in areas below the mid-shin around the foot and ankle where the skin is more sensitive. Although this generally settles, it can sometimes take a number of weeks or months. Continue to regularly massage the area with moisturiser to de-sensitise the areas.

Venous Thromboembolic (VTE)

Although procedural related VTE complications are rare after sclerotherapy, deep vein thrombosis (DVT) and pulmonary embolism (PE), can still occur in 0.5% to 1% of treatments. Continued mobilisation and use of compression hosiery serve to minimise VTE risk following sclerotherapy. However, it is still important to watch for symptoms of DVT which may cause significant swelling of your treated lower extremity, with throbbing, pain and warm/red discolouration. A PE could cause shortness of breath and/or chest pain.

Phlebitis

Any increase in tenderness or redness of the treated areas can sometimes represent a phlebitis (vein inflammation) which can occur soon after the procedure or even a few months later. It is recommended to take oral anti-inflammatory medication such as Ibuprofen 200mg 3 times a day (with meals to avoid irritation to your stomach) for 1-2 weeks and only if it is safe for you to take oral anti-inflammatory medication.

If you have a history of previous phlebitis, Professor O'Donnell will have already advised you to take oral Ibuprofen tablets 3 times a day for 2 weeks post-procedure as a preventative measure if it is safe for you to take Ibuprofen or an alternative oral anti-inflammatory tablet.

Occupational Duties & Travelling

Patients may return to work whenever they wish after sclerotherapy. Driving may be recommenced the following day after treatment. There are no specific precautions for short-haul flying after treatment but it is recommended to keep yourself well hydrated, mobilise gently on the flight and wear your compression stockings if flying in the first 2 weeks after your procedure. Long-haul flights (over 4 hours) are best avoided in the first 4-6 weeks after your procedure.

Sun Exposure

It is recommended to limit sun exposure in the first 6 weeks after your treatment as this can result in prolongation of procedural bruising and an increased risk of skin pigmentation. It is prudent to try and ensure that you avoid sun exposure to your treated lower extremities until 6 weeks after treatment or after any subsequent review consultations with Professor O'Donnell in case further treatment is administered during the same consultation.



Follow up

Professor O'Donnell will normally review you at his clinic approximately 3 months post-procedure to check on your progress and make sure your treatment outcomes are satisfactory. If you have any concerns, please contact your clinic.

Note:

It is ok to re-schedule your clinic appointments with Professor O'Donnell.



Clinic Opening Hours & Contact Information



Kingsbridge Private Hospital Belfast

811-815 Lisburn Road, Belfast, BT9 7GX

Daytime | 028 9066 7878
Out of Hours | 028 9068 8867



Kingsbridge Private Hospital North West Ballykelly

Church Hill House, Main Street, Ballykelly, BT49 9HS

Daytime | 028 9066 7878
Out of Hours | 028 7772 3506



Kingsbridge Diagnostic & Treatment Centre Belfast

Building One, Dataworks, King's Hall Life Sciences Park, Belfast, BT9 6GW

Daytime | 028 9590 9605
Out of Hours | 028 9068 8867



Kingsbridge Newry Clinic Newry

Windsor Avenue, Newry, BT34 1EG

Daytime | 028 3025 7708 / 028 9066 7878
Out of Hours | 028 9068 8867



We wish you a smooth and speedy recovery and look forward to continuing to provide you with the highest quality healthcare.

Professor Mark O'Donnell & the Varicose Vein Clinic Team
at Kingsbridge Private Hospital Group.



KINGSBRIDGE

Private Hospital Group



Kingsbridge Diagnostic & Treatment Centre Belfast

Building One, Dataworks, King's Hall Life Sciences Park,
Belfast, BT9 6GW

Kingsbridge Private Hospital Belfast

811-815 Lisburn Road,
Belfast, BT9 7GX

Kingsbridge Private Hospital North West

Church Hill House, Main Street,
Ballykelly, BT49 9HS

Kingsbridge Newry Clinic

Windsor Avenue,
Newry, B349 1EG



[kingsbridgeprivatehospital.com](https://www.kingsbridgeprivatehospital.com)